

## **Historic, archived document**

Do not assume content reflects current scientific knowledge, policies, or practices.


1884  
Ag 84  
#132



# U. S. DEPARTMENT OF AGRICULTURE

FARMERS' BULLETIN No. 1353 *rev.*  
*May 1935*

## CLOTHES MOTHS AND THEIR CONTROL



**C**OMplete elimination of clothes moths from dwellings and other buildings is difficult. These insects breed not only in wearing apparel, carpets, rugs, piano felts, and upholstered furniture, but in the woolen lint lodged in floor cracks and similar places.

Clothes moths can be kept under control, however, by persistent and intelligent use of the measures described in this bulletin.

Women's clubs and neighborhood organizations can greatly reduce expenses in work against moths by pooling their needs and purchasing in quantity from local dealers or manufacturing chemists the insecticides required. Naphthalene or paradichlorobenzene, two of the safest and best materials for protecting fabrics in storage against moth injury, can be obtained in this way.

# CLOTHES MOTHS AND THEIR CONTROL

By E. A. BACK, *principal entomologist, Division of Insects Affecting Man and Animals, Bureau of Entomology and Plant Quarantine*

## CONTENTS

	Page		Page
Clothes moths the chief pests of fabrics.....	1	Control of clothes moths—Continued.	
Harmless moth miller parent of destructive worm.....	1	Pyrethrum powder.....	16
Common species of clothes moths.....	2	Cold storage.....	16
The case-making clothes moth.....	2	Storage in vaults with fumigation.....	18
The webbing clothes moth.....	3	Fumigation of entire establishment with hydrocyanic acid gas or sulphur fumes.....	19
The tapestry moth.....	5	Carbon disulphide fumigation.....	21
Food of clothes moths.....	5	Carbon tetrachloride fumigation.....	24
Biology of clothes moths.....	7	Fumigation with ethylene dichloride-carbon tetrachloride mixture.....	24
The adult or moth miller.....	7	Fumigation with mixtures of ethylene oxide and methyl formate with carbon dioxide.....	25
The egg.....	8	Cedar-lined closets.....	25
The larva.....	9	Cedar chests.....	25
The pupa or chrysalis.....	10	Cedar products.....	27
Seasonal history.....	10	Dry heat.....	28
Excrement mistaken for eggs.....	12	Oil sprays.....	28
Control of clothes moths.....	12	Hot water.....	28
Ineffective measures.....	13	Laundry soap.....	29
Constant watchfulness.....	13	Gasoline.....	29
Mothproofing solutions.....	13	Treatment of cracks and hiding places with gasoline, benzene, or kerosene.....	29
Brushing, beating, and sunning.....	13		
Paper garment bags.....	13		
Careful wrapping in unbroken paper.....	14		
Moth crystals and moth balls.....	14		

## CLOTHES MOTHS THE CHIEF PESTS OF FABRICS

**C**LOTHES MOTHS are the prime offenders among fabric pests. The moths or "millers" most common about houses have a wing spread of about half an inch and are yellowish or buff colored. They very seldom fly directly to a light in a room and remain flitting about it, as do so many small and large moths that enter houses from outside. Moths that fly in large numbers about the home light are generally feeders upon outdoor vegetation and will not eat household fabrics. Their presence in the house is accidental—they have merely been lured inside by the lighted lamp.

The common clothes moths are usually seen flying in darkened corners and just beyond range of the brightest rays of the lamp. They prefer darkness. They are frightened when clothing and other objects are suddenly moved, and are then seen running rapidly or flying to conceal themselves in the creases of clothing, cracks, or other dark places.

## HARMLESS MOTH MILLER PARENT OF DESTRUCTIVE WORM

The moths, or millers as they are often called, have imperfectly developed mouth parts and could not feed upon fabrics if they desired. Their purpose in life is merely to lay eggs that develop into the worms or larvae, which alone can cause destruction. Nevertheless, every moth killed helps to reduce infestations by preventing more eggs being laid. The moths lay their eggs in and about clothing and other objects subject to attack. From the eggs are hatched

the larvae, or worms. When these become fully grown, they are whitish and about half an inch long. The larvae spin cocoons in which they transform to the pupa or chrysalis stage, and during this stage the insect changes to the adult, or moth. Thus each generation of clothes moths passes through four distinct stages, namely, the egg (fig. 14), the larva or worm (figs. 1, 2, 4, and 7), the pupa or chrysalis, and the adult or moth (figs. 1, 4, 6, and 7). Only the worm or larva of the clothes moth is capable of destroying fabrics.

#### COMMON SPECIES OF CLOTHES MOTHS

There are two very common species of clothes moths. They are the case-making clothes moth (*Tinea pellionella* L.) and the webbing clothes moth (*Tineola biselliella* Hummel). The tapestry moth (*Trichophaga tapetella* L.) is less often found, though it may become destructive.




FIGURE 1.—Case-making clothes moth: Above, adult; at right, larva; at left, larva in case. Enlarged. (Riley)

#### THE CASE-MAKING CLOTHES MOTH

The adult of the case-making clothes moth (*Tinea pellionella* L.) has a wing expanse of about half an inch. Its head and forewings are grayish yellow or buff, with

indistinct dark spots on the middle of the outer portions. The hind wings are white or grayish and silky.

The case-making clothes moth (fig. 1) is so named because the larva, for its protection, makes a portable case out of spun silk and fragments of the fabric upon which it feeds. These cases attain a length of from one-quarter to three-eighths of an inch and are shaped as illustrated in Figures 1 at left and 2, *a, c*. The larva almost never leaves its case. It withdraws completely into it when resting, but when feeding or moving from place to place it merely protrudes its head and first body segments, as illustrated in Figure 1, at left. The larva spins almost no web on the fabric upon which it feeds, and is more likely to crawl about restlessly, eating small holes here and there. (Fig. 2, *d*, and Fig. 3.)

As the larva grows it enlarges its case from time to time, both in length and in circumference. This enlarging process has been interestingly described as follows:

Without leaving its case the larva makes a slit halfway down one side and inserts a triangular gore of new material. A similar insertion is made on the opposite side, and the larva reverses itself without leaving the case and makes corresponding slits and additions in the other half. The case is lengthened by successive additions to either end. Exteriorly, the case appears to be a matted mass of small particles of wool; interiorly, it is lined with soft whitish silk. By transferring the larva from time to time to fabrics of different colors the case may be made to assume as varied a pattern as the experimenter desires,

and will illustrate, in its coloring, the peculiar method of making the enlargements and additions described.<sup>1</sup>

On reaching full growth the larva attaches its case by silken threads to the material upon which it has been feeding, or it may crawl, carrying its case, to another part of the room. In either instance, after attaching the case, it seals it with silk, thus making a pupal chamber wherein it transforms from the helpless pupa or chrysalis to the pretty and active moth.

#### THE WEBBING CLOTHES MOTH

The adult (fig. 4, upper half) of the webbing clothes moth (*Tineola biselliella* Hummel) is of about the same size as that of the case-making species, but its color is uniformly pale buff without spots. The moths of both species vary considerably in size; they seldom have a wing expanse of over half an inch, and frequently are very much smaller.

The webbing clothes moth is the most abundant and injurious clothes moth. During the last few years, also, practically all instances of severe clothes-moth damage reported by manufacturing plants throughout the North, including Chicago, Boston, and New York, have been associated with this species. For years it has been the prevailing species in certain houses in western Massachusetts and southeastern Connecticut; hence it can not be said that this webbing clothes moth is typically southern in distribution, although it is often called the "southern" clothes moth.

The larva (fig. 4) of the webbing clothes moth resembles that of the case-making moth. Unlike the latter, however, it makes no portable case, but spins silky transparent tubes or tunnels wherever it goes. (Figs. 5 and 13.) Often it spins a cobwebby mass of




FIGURE 2.—Case-making clothes moth: *a*, Three cases nearly twice natural size; *b*, larva removed from cases, and pellets of excrement; *c*, cases attached to damaged feather, about one and one-third times natural size; *d*, characteristic damage by larva to cloth

<sup>1</sup> C. L. Marlatt, in Bulletin 4, new series, Division of Entomology, U. S. Department of Agriculture, p. 65. 1896.

silken threads, apparently at random, as it crawls about fabrics, which gives to its food an unsightly appearance. The larvae may be quite restless, and often may be seen crawling over fabrics, or




FIGURE 3.—Section of man's coat damaged by case-making clothes moth. Note large eaten area at lower corner and along lower edge and small holes eaten at arrows

upon floors beneath upholstered furniture from which they have dropped. They attain a length of about half an inch and are white. Figure 4 gives a good idea of the relative size of the larvae and moths; the buttons shown are nine-sixteenths of an inch in diameter. When the larva becomes full grown, it spins a cocoon of silk, inter-




FIGURE 4.—Webbing clothes moth : Adults or moths above and larvae or worms below. Buttons are nine-sixteenths of an inch in diameter

mixed with bits of fabric and excrement, which is quite distinct from the case (fig. 2, *a*, *c*) made by the case-making clothes moth. Within this cocoon the transformation from the larva to the adult

takes place, and, as in the case of the other species, the chrysalis works its way partly out of the cocoon as the moth is about to emerge.

#### THE TAPESTRY MOTH

The tapestry moth (*Trichophaga tapetzella* L.) (fig. 6) is not so common in the United States as the two foregoing clothes moths. It is larger than either the case-making or the webbing clothes moth, having a wing expanse of about three-fourths of an inch, and is more strikingly colored. Its head and the basal third of its forewings are black, while the outer two-thirds of the wings are creamy white, though more or less obscured on the middle with gray. The hind wings are uniformly pale gray.




FIGURE 5.—Work of the webbing clothes moth in piece of cloth with heavy nap. Note long, narrow tube in which the larva secretes itself and from which it feeds upon the cloth. To the right of the tube is shown an area over which the larva has eaten off the nap, leaving the warp exposed.

The tapestry moth is said to prefer coarser and heavier fabrics than the other two species of moths. It is more likely to occur in carpets, horse blankets, tapestries, felting, furs, and skins. The larvae construct burrows or silk-lined galleries in all directions throughout the infested material, and often cause quite as much injury by these as by the amount of the fabric that is actually eaten. The tapestry moth has been reported as destructive to wall paper.

#### FOOD OF CLOTHES MOTHS

Clothes moths feed upon wool, fur, hair, feathers, and all fabrics (figs. 7, 8, and 9) manufactured from them. They also relish dried



animal matter such as dead insects, including the dead of their own species, and untreated skins, beef meal, casein, etc. In experimental work clothes moths have thrived best upon a diet of bristles, hair, feathers, fur, or raw wool, and not so well upon ordinary woolen cloth used in the making of wearing apparel. Notwithstanding this, the damage they cause to wearing apparel has given them a well-




FIGURE 6.—Tapestry moth: Adult. Enlarged. (Riley)

deserved reputation as pests. There may be some satisfaction, however, in knowing that if the death rate among clothes moths feeding upon dyed woolen fabrics was as low as among those feeding upon raw wool, and their growth as rapid as when they feed upon bristles, fur (figs. 10 and 11), and hair,

their capacity for destruction would be increased enormously. When a campaign is waged against clothes moths in the home, it must be remembered that they may be found feeding upon carpets and rugs (figs. 7, 8, and 9), stuffing and covering of upholstered furniture, tapestries, stuffed animals or birds, dust brushes (fig. 12), shaving brushes (fig. 13), the felts in pianos, and a long array of articles often little thought of as being fed upon by clothes moths. Woolen lint, and hair from pet animals which




FIGURE 7.—Clothes-moth injury to carpet: Side view of carpet showing moth, with eggs on tuft, to left, larva eating at base of tuft to right, and cocoon with end cut to expose pupa within. Note larva crawling over tuft at extreme right. Greatly enlarged

has become lodged in floor cracks, behind baseboards, and elsewhere can keep moths alive in otherwise scrupulously clean houses. Clothes moths almost never injure household fabrics or wearing apparel in daily or weekly use, or articles that are brushed, cleaned, or beaten once or twice a month. Furniture, on the other hand, may be in constant use and still be badly infested.

## BIOLOGY OF CLOTHES MOTHS

It is important in a fight against clothes moths to know something about their biology. Each generation is called a life cycle because the insect keeps passing through a successive round of stages, consisting of the egg, the larva, the pupa, and the moth. The fol-



FIGURE 8.—Clothes-moth injury to rugs. When the injury illustrated above has progressed sufficiently far, whole patches of pile are cut off by the larvae and the warp is left exposed as here shown. Sometimes larvae feed on the underside of carpets and rugs, but they cut the pile there just as in feeding as indicated in Figure 7

lowing facts concerning these stages of the life cycle relate particularly to the webbing clothes moth, but must be taken into consideration in any intelligent campaign against clothes moths of other species as well.

## THE ADULT OR MOTH MILLER

The moth, or "miller," which is the adult, seldom lives as long as one month. Moths are more likely to die between the seventh and




FIGURE 9.—Clothes-moth infestation in rugs and carpets can be detected by bending the fabrics so that the space between the tufts of pile can be examined. Moth webbing and cocoons (as here shown) are thus readily exposed

fourteenth days after they emerge from the pupa. Apparently they take no nourishment, and the female moths begin laying eggs before they are fully 1 day old and usually lay eggs each subsequent day of their lives. When a moth stops laying eggs it is a sign that she will die within a day or two. Few moths have been known to lay over 100 or 150 eggs, and of this number at least one-half

are laid during the first few days of adult life. As many as 59 eggs have been laid during 24 hours by one female moth, but this is an unusual number.

#### THE EGG

The eggs are laid singly or in groups of as many as 25, either loosely upon or between folds of fabrics. In loosely woven goods, such as yarns, carpets, and rugs with heavy nap, or heavy woolen clothes, the eggs are placed down in the meshes of the fabric, where they are held by the fine threads, as illustrated in Figure 14. On certain closely woven serges they are laid so loosely upon the sur-




FIGURE 10.—Aviator's helmet slightly damaged by webbing clothes moth larvae. Compare with Figure 11, which shows complete destruction of fur

face that they can be shaken off. In furs, the female moth may burrow down among the hairs to the skin itself in order to find a sheltered place for her eggs. The eggs are not as large as the head of a common pin, but, small as they are, their white color makes it possible to see them without the aid of a magnifying glass, if they are laid upon a dark or black substance. They are very fragile and easily crushed.

Clothes-moth eggs hatch most readily in warm summer weather in from four to eight days. During colder weather hatching may not take place for three weeks, but in steam-heated buildings it does not require a much longer time than it does in warm summer weather.

## THE LARVA

Of all the stages, the larva or worm stage—the only stage in which clothes moths damage fabrics—is the most susceptible to outside influences. The nature of its food, the temperature, and the humidity have pronounced effects upon its growth. Sometimes, for no apparent reason, the larva may pass into a period of dormancy lasting as long as 8 to 24 months, during which it will neither feed nor move about appreciably, only later to become active, feed, and continue its growth.



FIGURE 11.—Aviator's helmet with fur completely eaten away by larvae of webbing clothes moth. Note cocoons of moth on upper portion, and that the larvae have even eaten the leather on central lower portion

When first hatched the larva is white, as when full-grown, and is scarcely one-sixteenth of an inch long. When young it is so transparent that the color of its food shows through its alimentary canal as a line extending along the center of the body. When full-grown it may be half an inch long, although its size will vary with its environment and food supply. It is not possible to judge the age of a larva by its size. Some larvae at 15 months of age are very much smaller than others only 2 months old. The shortest definite record for larval development known is about 40 days at 80° F.

Many larvae hatching during the summer do not become fully grown and mature as moths until the following March, April, and May. There even is a considerable variation in the length of time larvae of the same brood require for development. Larvae hatched

on the same day may require as few as 15 weeks or as long as 4 years. Much depends upon their food and environment.

#### THE PUPA OR CHRYSALIS

The pupa or chrysalis stage requires during warm summer weather about 8 to 10 days. During the winter months from January to March, in a steam-heated building, clothes moths remained in the pupa stage three to four weeks.




FIGURE 12.—End view of common bristle dust brush, showing sandlike pellets of frass and damaged bristles. Clothes moths can feed upon all articles made of bristles

#### SEASONAL HISTORY

While moths may be on the wing in steam-heated buildings during almost any month, they are present in greatest abundance the country over from May to July and during September and October. In steam-heated houses moths of the webbing species have been observed as early as late December, January, and February. In northern localities, where only the case-making moth is present, and in houses not continuously heated, the moths are single-brooded and appear during late May, June, and July. The late appearance of moths in such northern houses is greatly to the advantage of the housewife, for by June winter clothing can be stored where it is protected from moth eggs.




FIGURE 13.—End view of a shaving brush, showing a well-developed larval tube of the southern or webbing clothes moth




FIGURE 14.—Eggs of webbing clothes moth: At *a* and *c*, laid among the threads of a loosely woven cloth; at *b*, laid upon a closely woven broadcloth. Greatly magnified

While it is true that wearing apparel, if in daily or weekly use, is not damaged by clothes moths, the occasional appearance of moths in steam-heated houses in late winter makes it clear that moth eggs may be laid on susceptible articles at any time they are not in use, and that if these are laid away without precautionary measures being taken to protect them they may be damaged in storage.

#### EXCREMENT MISTAKEN FOR EGGS

When articles very badly damaged by clothes moths are unfolded there are always seen tiny sandlike particles, hard and gritty, which vary in color according to the color of the food upon which the moth worms have been feeding. These particles are pellets of larval excrement and not eggs. A single moth larva may void about 700 of these pellets during its growth. Fortunately eggs are never present in such numbers (pp. 7 and 8).

#### CONTROL OF CLOTHES MOTHS

There are effective methods, discussed in this bulletin, for controlling clothes moths and reducing to a minimum the losses they cause. The fumigation of houses is not always feasible or necessary. Thoroughly cleaning, brushing, and, if possible, sunning fabrics and immediately storing them with naphthalene or with paradichlorobenzene is an effective and inexpensive precaution.

#### INEFFECTIVE MEASURES

No repellents for clothes moths are known. The belief that the mere smell of certain substances keeps or drives moths away has been proved to be incorrect. In Department experiments a room was equipped with a number of closets, the doors of which were left slightly ajar, and adult moths set free in the center of the room entered the closets containing the odor of naphthalene, paradichlorobenzene, cedar, pine, or camphor, as soon as the closets without odors. Clothes hung in the closets containing small quantities of the above-mentioned substances were as badly damaged by the larvae as were the clothes in the other closets. The mere *smell* of a chemical does not drive moths out of clothing nor does it keep them out.

The following substances often recommended for clothes moth control are worthless for this purpose:

Tobacco extracts containing nicotine and tobacco powder when used at reasonable strengths.	Quassia chips (dusted).
Lavender flowers (scattered on).	Borax (dusted).
Cayenne pepper.	Colocynth pulp (dusted).
Allspice (dusted).	Eucalyptus leaves (dusted).
Angelica root (dusted).	Helebores, white (dusted).
Black pepper.	Formaldehyde (sprayed 1 to 10).
Pyrethrum stems (dusted).	Sodium bicarbonate (dusted).
Lime, air-slaked (dusted).	Sodium carbonate (dusted).
Powdered sulphur (dusted).	Lead carbonate (dusted).
Salt (dusted).	Lead oxide (dusted).
	Red cedar leaves (dried and placed in clothing).

### CONSTANT WATCHFULNESS

Naturally, constant watchfulness must be the watchword for successful control. No treatment to kill clothes moths already in fabrics will have any lasting effect in keeping other clothes moths from later infesting these fabrics if left exposed about the house. No practical known treatment will render wearing apparel absolutely immune to attack.

### MOTHPROOFING SOLUTIONS

Numerous mothproofing solutions for which much is claimed by manufacturers are available. In many instances guarantees are issued to the effect that one application will render the treated fabric immune to moth attack for the life of the garment, or until the fabric is washed or drycleaned, depending upon whether the solution is removable by washing in water or by drycleaning fluids. From laboratory tests and the experience of the public, it would seem that no mothproofing solution now available, is capable of imparting a permanent or absolute mothproofness. It would seem, however, that some of the better fluoride, cinchona alkaloid, and rotenone solutions, if so used as to thoroughly impregnate the treated fabric, do impart a moth resistance that is of very considerable value, and when a person applies such a treatment he gets the best protection of this nature available.

Solutions depending for their effectiveness on an arsenic content are not recommended by this Department because of the danger of skin troubles developing as a result of contact with arsenic-treated fabrics or furs.

### BRUSHING, BEATING, AND SUNNING

Too much emphasis cannot be placed on the value of frequent brushing, beating, sunning, and cleaning of articles subject to clothes-moth attack, as the eggs of clothes moths are delicate they can usually be crushed or dislodged by brushing and beating. Pay special attention to crevices, seams, and pockets. Sunning is a valuable aid in control work. In articles laid away moths are much more likely to concentrate upon soiled spots if these have not been removed. Brushing and beating usually remove or kill the larvae or worms. If clothing is thoroughly brushed every two weeks, it is doubtful if moths can affect it seriously. Carpets and rugs cleaned by strong vacuum cleaners and thoroughly brushed on both sides, or electrically cleaned, are freed from infestation if the work is properly done. Clothing dry-cleaned is also freed from infestation by the process. Once freed from infestation, clothes should be protected, if they are not to be used during the warm season, by being wrapped in paper or placed in tight chests or treated closets.

### PAPER GARMENT BAGS

The public is familiar with "mothproof" paper bags of various types that appear upon the market each summer. These paper bags for the storage of suits and coats are made of heavy paper, paper impregnated with tar, or other substances for which much is claimed.



These bags if not torn and if properly used are excellent for keeping moths from reaching clothing. They are of no value in killing moths that may be in the clothing when it is placed in the bag, and those made of plain paper are just as good as those containing tar or cedar and pine oils.

The value of such bags as protectors against infestation ceases largely as soon as they are torn, even slightly, or if they are left unsealed. Frequently clothing on coat hangers is placed in bags with the hook of the hanger protruding from the bag so that the clothing and bag may be hung in the closet. Unless special attention is given to this method of hanging, the paper bag is not carefully sealed about the hanger, and spaces of one-fourth inch or more are left open. Moths have no difficulty in crawling into such openings, and if they find them the bags are useless.

#### CAREFUL WRAPPING IN UNBROKEN PAPER

*Ordinary firm wrapping paper is satisfactory, or several thicknesses of newspapers will serve when one is wrapping garments or other fabrics.* Clothes moths do not eat into paper to reach clothing. For this reason, if woolens and other fabrics subject to moth attack are cleaned and freed from moths by any of the methods mentioned above, they will remain safe if wrapped at once and tightly in several thicknesses of firm wrapping paper or in newspapers. There is a general belief that the print on newspapers keeps moths away. This is not true. Any unbroken paper of moderate thickness will protect clothing. Care must be exercised to bend back the folds of the paper upon themselves along the edges and at the ends of the bundles so that the moths cannot crawl into them. Hats and other articles that might be damaged by tight wrapping can be protected by being placed in ordinary unbroken hat boxes, cardboard boxes, etc., and the covers of the boxes sealed with gummed paper.

After clothing has been made into bundles or sealed in boxes, these bundles and boxes may be left exposed in garrets or on storeroom shelves without danger of infestation from without. If flake naphthalene is wrapped among the folds of clothing made into bundles, any small larvae or eggs that may have got into the cloth before it was wrapped will be killed.

#### MOTH CRYSTALS AND MOTH BALLS

The crystalline substances commonly used in moth control are naphthalene, paradichlorobenzene, and gum camphor. These can be purchased at drug or department stores and are of the greatest value and completely effective only if used in sufficient quantity and in tightly closed containers such as trunks, closets, or even rooms.

In evaporating slowly the crystals form a vapor or gas which is heavier than air and tends to sink to the bottom of the container. These vapors will even pass through paper if it is untreated. The mere smell will not protect garments (p. 12), and if the trunk or closet is not airtight and the vapors easily leak out, no practical protection is offered. But if the lid, or the closet door, shuts tightly against a felt strip or gasket and is kept shut most of the time, and

the closet is tightly constructed otherwise, genuine protection is given, for the vapors remain in sufficient concentration over a long enough time to prevent the moth worms from feeding and ultimately to kill them. The secret of success is to use only tight containers. If they are not as tight as they should be for the best results, make them as tight as possible by pasting strips of heavy paper over cracks or using crack fillers and employ a greater quantity of the chemical so that the vapors that escape will be replaced by others formed during the slow evaporation of the crystals.

#### NAPHTHALENE

Naphthalene is well known in the form of flakes or moth balls. If in good condition, it is one of the safest and best materials for protecting garments, rugs, etc., against moths. It is cheap and can be depended on if used in sufficient quantity; for long periods of storage it is as good as any crystalline substance for protection against moths, though its odor is not pleasant.

Naphthalene flakes cost from 6 to 20 cents a pound. They should be used at the rate of about 1 pound to each 6 to 10 cubic feet of space, which is the size of the average trunk. Experiments have proved that 2 or 3 ounces of flake naphthalene, or 8 ounces in the form of moth balls, will kill all stages of the moths if confined in a *thoroughly tight* container having a capacity of 5 cubic feet. In the opinion of the writer it is safer to use at least 1 pound of flake naphthalene scattered between the folds of clothing in a trunk of average size. You can thus be certain of killing the young and old moth worms, moth eggs, and the moths themselves if they happen to be present. A moth ball here and there will do no good. When large closets or rooms are used for storage no moth can cause injury if enough flake naphthalene is used to cause the eyes and nose to smart when one enters.

If a good grade of naphthalene is used there will be no sooty deposit when the crystals evaporate, but as a guard against impurities that may stain delicate fabrics it is wise to scatter the flakes between folds of old linen or tissue paper. In treating a closet, scatter the crystals well over shelf and floor before closing the door.

One advantage in using naphthalene compressed into moth balls or cakes is that these can be put in bags made of cheesecloth and hung from hooks in the closet. These balls or cakes evaporate slowly, thus lasting longer, but for quick results the flake form is better as the evaporation is much more rapid.

The vapors escaping from closets through cracks or when the door is opened will not be strong enough to harm a person outside.

#### PARADICHLOROBENZENE

Paradichlorobenzene, like naphthalene, can be had in the form of flakes or crystals, or compressed into forms variously colored and variously perfumed to mask the true odor, which is, to some, less objectionable than that of naphthalene. Paradichlorobenzene enjoys popularity because its odor is more pleasing to some and is less persistent, and because a greater effort has been made to merchandise

it in an attractive manner. For this the public pays an added price. In pound tins it sells for about 50 or 60 cents.

As in the case of naphthalene, it is best to scatter the crystals between layers of tissue paper distributed throughout the clothing in the trunk or chest. Paradichlorobenzene is used in the same manner as naphthalene, and is more quickly effective in stopping the feeding of the clothes moths than is naphthalene, but this fact is of little practical value if one is storing goods for the summer months, because for such a period the cheaper naphthalene will protect just as well, as is proved by its general use in the warehousing industries.

In a tight container one-half ounce of paradichlorobenzene for each 100 cubic feet of space has been reported to kill all stages of the moths in 31 days. Two ounces for the same space stops feeding quickly. But because no two household storages are equally tight unless specially built and because garments to be protected are always relatively valuable, the writer advises for complete safety the use of 1 pound of the crystals to each trunk of average size, although some advocate much less. Unless you know that your container is really tight it would be much better to use 1 pound of flake naphthalene than 2 ounces of paradichlorobenzene, as the pound of the former would last a considerable time and continue to give off vapors to replace those lost by leakage.

*Remember that neither paradichlorobenzene nor naphthalene protect your garments longer than the crystals last.* Unless they have been used in quantities sufficient to kill and in containers so tight that new moths cannot get in, they must be renewed after evaporation. It is for this reason that the heavier dosages are recommended.

#### CAMPHOR

Gum camphor used in the manner recommended for naphthalene and paradichlorobenzene will kill all stages of the clothes moth. If of good quality and used at the rate of from one-half to one pound to each 5 cubic feet of space in tight containers, gum camphor will protect. To be most effective the camphor should be broken into small pieces and used in tight containers which will confine the fumes. While gum camphor does kill some eggs and larvae when sprinkled in small pieces upon infested cloth in open containers such as bureau drawers, pockets in clothing, etc., it is not to be depended upon except where the fumes given off by evaporation can be closely confined with the clothing to be protected.

#### PYRETHRUM POWDER

Pyrethrum powder, if fresh, will kill clothes moth larvae. Clothing should be thoroughly dusted with the powder and placed in a tight container such as a chest or trunk, or wrapped well in unbroken paper. Pyrethrum powder soon loses its protective value on exposure to air, hence it is not considered particularly effective for long storage and is inferior to naphthalene or paradichlorobenzene. It is not always easy to obtain fresh pyrethrum powder at stores.

#### COLD STORAGE

Protection against injury by fabric pests of all kinds is assured by proper cold storage. Once fabrics are in cold storage, no injury can

take place. Cold storage (fig. 15) is depended on for absolute protection by dealers in carpets, furs, and other valuable articles such as stuffed animal heads, blankets, automobile robes, curtains, and upholstered furniture. Cold-storage facilities have increased with public demand until today no city or large town is without storage equipment. A tour during the summer months through the cold-storage rooms of large department stores and storage concerns with their wealth of furs and woollens cannot but impress one with the well-grounded faith of the public in this means of protection.

Articles will be protected from injury in storage at temperatures ranging from  $40^{\circ}$  to  $42^{\circ}$  F. A number of years ago a manager of a large storage-warehouse company in Washington, D. C., conducted certain experiments at the instance of the Chief of the Bureau of Entomology, with the result that it was found that larvae of the



FIGURE 15.—View in cold-storage room for rugs and carpets in a modern fireproof storage warehouse. Rugs and all other articles subject to attack by fabric pests are absolutely protected while in cold storage.

webbing clothes moth and of the black carpet beetle can withstand for a considerable time a temperature of  $18^{\circ}$ . It has been discovered that it is not so much the cold that kills. It is the sudden change from a cold to a warmer temperature and back to a cold temperature that most quickly results fatally. Thus it was learned that if articles infested with clothes moths were refrigerated at  $18^{\circ}$  for several days, then suddenly exposed for a short time to  $50^{\circ}$ , and then returned to  $18^{\circ}$ , and finally held permanently at about  $40^{\circ}$ , all moth life in them would be killed.

If storage concerns intend to destroy the clothes moths in articles intrusted to them, as well as to protect them from injury during the period of storage, it is recommended that the articles be exposed to

two or three changes of temperature as noted above before they are placed permanently at 40° to 42° F. The maintenance of a temperature lower than 40° to 42° is needless and a wasteful expense.

If the storage company merely guarantees to protect articles during the period of storage, it is quite sufficient to maintain them at a temperature of about 40° to 45° F. Companies should advise patrons that clothes-moth larvae can withstand storage at temperatures ranging from 24° to 45° for longer periods than that for which the majority of articles are refrigerated. Thus well-grown larvae of the webbing clothes moth in fur and wool held in commercial cold storage at a temperature said to fluctuate between 24° and 48°, but held mostly at about 40°, were found by the writer to be alive after storage for 6, 8, 10, 11, and 12 months. Larvae in fur helmets placed in storage during February, removed the following December, and held in a steam-heated building, transformed in considerable numbers into active adults by the middle of January. These adults showed no effects of the refrigeration of the larvae from which they had developed, but laid many eggs that hatched normally. Refrigeration for 6 months had no noticeable effect upon the larvae, except to hold them inactive and incapable of causing injury. While larvae refrigerated for 10 to 12 months matured, as above stated, into moths, a large percentage of these moths, though active soon after removal from storage, died after being subjected to warmer temperature. These facts are important, for they explain why patrons have at times found living, active, and robust larvae in articles several days after they had been removed from the cold storage in which they had been held 4 or 5 months. If a fairly even temperature has been maintained, such a discovery is possible and is no reflection upon the storage concern. If articles are cleaned thoroughly before being placed in storage of any sort, as they always should be, the likelihood that they will contain well-grown larvae is very slight.

#### STORAGE IN VAULTS WITH FUMIGATION

During the last few years an increasing number of storage firms have installed burglarproof and fireproof storage vaults in which furs, coats, rugs, furniture, and other articles can be stored and absolutely protected from moths by fumigation. This method of storage combines tight storage and fumigation. Such treatment has long been recommended by this Department. When intelligently followed, this method of protecting articles against moth ravages is excellent and is the equal of any other. Because the cost of installing and operating this type of storage is moderate and because the results are satisfactory, it has been adopted by some of the largest department stores and storage concerns for storing furs and other articles during the summer. (Fig. 16.)

Operators of this type of storage and curators of furs at the National Museum have found that the luster and natural oils of tanned furs are not affected by such storage, although statements to the contrary are often heard. The life of the fur is prolonged by this method as it is by any other storage method approved by this Department, because the fur is protected from destructive insects, light, dust, and excessive humidity.

FUMIGATION OF ENTIRE ESTABLISHMENT WITH HYDROCYANIC ACID GAS  
OR SULPHUR FUMES

In large houses with many carpets or rugs, much upholstered furniture, and other fabric furnishings, and in tourist hotels and houses used only during the winter months, but closed through



FIGURE 16.—View in modern storage room where furs and other garments are protected perfectly from moth attack by applying the principle of tight storage and fumigation.

the long, hot summer season, fabric pests may become very abundant and so generally established that they cannot be reached easily by the more local treatments recommended farther on. In

such establishments insects may not only be secreted in the affected furnishings, but may be upon the walls, in floor cracks, behind mopboards, and in other places where they have crawled for protection or for transformation. The removal of furnishings and their treatment outside the establishment, or in special rooms, will remove at the same time most of the infestation; yet enough insects may be left behind to restock the furnishings when they are returned after treatment. For this reason, any control measure that can be applied to the house or hotel with the furnishings left as they are normally has its advantages. Exception should be made in the case of chests or other tightly packed containers. Clothing, blankets, and other contents of such containers should be removed and hung up in rooms or variously spread out over furniture, etc., so that fumes can more readily reach the larvae in them. This need not be done if the containers are treated as separate units as hereinafter suggested. It may be well to raise rugs or carpets so that the gas can more easily reach the underside, though this is not always necessary or feasible. The tops of pianos and organs (upon the felts of which the moths and carpet beetles may be feeding) should be raised, and all closet doors, bureau drawers, trunks, etc., should be left partially open.

Fumigation by either of two methods can be used for the treatment of houses or hotels as a single unit—namely, with hydrocyanic acid gas or with sulphur fumes. *Formaldehyde fumigation is worthless in controlling clothes moths or other fabric pests, notwithstanding the general belief regarding it.* Formaldehyde is a good disinfectant for use after contagious diseases, but should never be employed in controlling fabric pests.

#### HYDROCYANIC ACID GAS

Fumigation with hydrocyanic acid gas is the best known method of controlling fabric pests as well as other household pests, including the bedbug. It kills speedily and effectively. The gas is formed by the union of sulphuric acid, sodium cyanide or potassium cyanide, and water, or given off by certain other cyanide products. It is colorless, lighter than air, with a distinct odor, and when mixed with air in the proportion occurring during fumigation is noninflammable. It is not injurious to the most delicate fabric or dyes, or to any house furnishing, and does not tarnish silver, brass, or other household metals. When the house is ventilated, the gas quickly escapes skyward, and houses usually can be ventilated so that they may be occupied within 1 or 2 hours, often within 30 minutes. Sleeping quarters at hospitals and private homes can be fumigated during the day and ventilated so that they can be safely occupied the same night.

*Hydrocyanic acid gas is deadly to human beings* if breathed in concentrated form and should be used only by an intelligent, careful, and well-informed person who understands the element of danger as well as the excellent features of the gas. Hydrocyanic acid gas should not be used in an apartment house, office building, or city block of houses unless neighboring parties are notified and are ready to vacate if necessary during the period of fumigation.

Hydrocyanic acid gas fumigation costs approximately 35 cents to \$1 for each 1,000 cubic feet of space fumigated. The house may be left exposed to the action of the gas for from 4 to 24 hours. Hydrocyanic acid gas fumigation is being employed in flour mills, warehouses, leading hospitals, schools, and colleges, as well as in private homes. For information on the method of fumigating with hydrocyanic acid gas see Farmers' Bulletin No. 1670.

#### SULPHUR FUMES

Fumigation with sulphur fumes is a remedy well known to the public. Sulphur fumes will kill clothes moths, but must be used at the rate of about 13½ ounces of sulphur to each 1,000 cubic feet of space. Fumigation should continue for a period ranging from 18 to 24 hours. *Remember* that there is some danger of fire in the careless use of sulphur candles; also that sulphur fumes may have a bleaching effect upon delicate fabrics and wall paper, and that they tarnish metals. The greater the atmospheric moisture the greater the bleaching effect. The Department of Agriculture does not recommend the use of sulphur fumes in homes equipped with valuable furnishings. Very often, however, the use of sulphur candles offers a most welcome method of moth control because of the availability of sulphur candles at drug stores. Naturally where fumes are used, rooms or houses must be tightly closed and more rather than fewer of the sulphur candles used. Failure with sulphur candles often is the result of dependence upon one or two improperly burned candles to do the work of the recommended 13½ ounces of sulphur for each 1,000 cubic feet of space to be fumigated. Directions for burning sulphur candles usually are supplied by the manufacturer or the dealer.

#### CARBON DISULPHIDE FUMIGATION

Carbon disulphide is an excellent fumigant for single rooms, closets (fig. 17), trunks (fig. 18), chests, and other tight containers. It is sometimes called "high-life", and can be purchased at drug stores or from wholesale chemical firms as a liquid put up in tin cans having a capacity of 1, 2, 5, 10, or more pounds at a cost of from about 6 to 35 cents a pound, according to the quantity used. It is a colorless<sup>2</sup> liquid that looks like water, but weighs about 10 pounds to the gallon. When exposed to the air it evaporates quickly, producing a foul-smelling gas about two and three-fourths times as heavy as air. While the liquid is not explosive or inflammable, the gas formed upon evaporation is. Because of this inflammable nature of the gas it must be kept away from fire in any form. The gas can be detected easily by its foul odor, but this odor disappears after thorough ventilation of rooms and fumigated articles. If used carefully and intelligently as directed and in the proper amounts, carbon disulphide speedily and surely kills fabric pests in tight containers. It is not recommended for the fumigation of entire buildings

<sup>2</sup> The ordinary commercial carbon disulphide has a decidedly yellowish color, owing to the excess of sulphur. This is a satisfactory fumigant, however, if too much of the impurities is not present.





FIGURE 17.—Clothing in closets can be freed of clothes moths if fumigated as illustrated with either carbon disulphide or carbon tetrachloride or the ethylene dichloride-carbon tetrachloride mixture. Since the gas formed by evaporation of these three liquids is heavier than air, pour the liquid into a shallow dish placed on the shelf of the closet. (Note white enamel dish on shelf.) As soon as liquid has been poured into dish shut door and seal cracks about door with gummed paper, or paste on strips of firm paper using a flour paste. Sealing the door prevents the gas from escaping rapidly.

because of the fire hazard when so great a volume of gas is formed, although experienced persons can use it safely for the fumigation of detached houses if they can control surrounding conditions.

When used in small amounts about houses in accordance with directions, there is but little more danger than in the use of gasoline or benzene, with which the public is more familiar. For complete information regarding carbon disulphide and its use, write to the United States Department of Agriculture for Farmers' Bulletin 799.



FIGURE 18.—Trunk ready for treatment with carbon disulphide, carbon tetrachloride, or the ethylene dichloride-carbon tetrachloride mixture. On the clothing are shown two saucers. In the one to the left stands a 1-pound tin can of carbon disulphide; in the one to the right a glass bottle filled with the colorless carbon tetrachloride. Any one of the three fumigants mentioned may be used. Pour the required quantity of the one to be used into the saucer and close and seal the trunk. The liquid evaporates, forming a gas heavier than air that sinks down into the clothing and kills clothes moths and other insects.

Fumigation with carbon disulphide is not effective at temperatures below 65° F. Better results follow where the temperature is above 70°. In rooms and closets or in well-built trunks the evaporation from 4 to 6 pounds of the liquid to each 1,000 cubic feet of space should kill all moths and their larvae. If the walls are plastered, or plastered and papered, the floors tight, and the doors sealed after fumigation begins, 4 pounds of carbon disulphide should prove sufficient to kill fabric pests. More liquid, even as much as

20 pounds per 1,000 cubic feet of space, may be required in more loosely constructed containers. Assuming that 4 pounds of carbon disulphide will be used to each 1,000 cubic feet of space, the following rough calculations may be useful in determining how much liquid is needed for various containers:

#### QUANTITIES OF CARBON DISULPHIDE NEEDED FOR FUMIGATION

Space to be fumigated	Quantity needed
Room, 12 by 10 by 8 feet-----	4 pounds or $3\frac{1}{4}$ pints.
Closet, 3 by 8 by 8 feet-----	$1\frac{1}{2}$ pints.
Closet, 2 by 5 by 7 feet-----	$\frac{3}{4}$ cup.
Closet, 4 by 5 by 7 feet-----	$1\frac{1}{2}$ cups.
Trunk, 21 by 20 by 42 inches-----	3 tablespoonfuls.
Box, 1 by 2 by 2 feet-----	1 tablespoonful.

Note that the method of application, as indicated in Figures 17 and 18, takes into consideration the fact that the gas formed is about two and three-fourths times as heavy as air. The liquid must therefore be exposed on top of the article to be fumigated.

#### CARBON TETRACHLORIDE FUMIGATION

Carbon tetrachloride is a good fumigant for single rooms, closets, trunks, chests, and other tight containers. It is a thin, transparent colorless liquid, in appearance similar to water, that evaporates on exposure to air, forming a gas with a pungent, aromatic odor. It is like carbon disulphide in that its gas is heavier than air. When carbon tetrachloride is used at the same rate as carbon disulphide, it is about one-half or one-third as effective. It has the great advantage over carbon disulphide that its gas is neither explosive nor inflammable; hence there is no fire risk in its use. Carbon tetrachloride is purchased at drug stores or from chemical firms in cans containing 1 or more pounds of liquid, and costs from  $10\frac{1}{2}$  to 30 cents a pound, according to the quantity purchased. It should be used at the rate of 8 to 12 pounds, or more, per 1,000 cubic feet of space if containers are very tight. When the temperature is  $70^{\circ}$  F. or above, good results in killing clothes moths should follow if the quantities given above for carbon disulphide for the respective spaces are trebled. Since the gas is heavier than air, the liquid should be exposed in shallow dishes at the top of the container to be fumigated. (Figs. 17 and 18.)

#### FUMIGATION WITH ETHYLENE DICHLORIDE-CARBON TETRACHLORIDE MIXTURE

Ethylene dichloride-carbon tetrachloride mixture consists of 3 parts by volume of ethylene dichloride and 1 part by volume of carbon tetrachloride. It is used in the same manner as are carbon disulphide and carbon tetrachloride. Like these, the gas produced by the evaporation of the liquid is heavier than air. Like carbon tetrachloride, the gas is noninflammable and nonexplosive and is not dangerous to human life in small quantities. It is about five times as toxic as carbon tetrachloride alone, but should be used at the rate of 5 quarts, or 14 pounds, per 1,000 cubic feet of space to be fumigated. Best results are obtained when the temperature is above

70° F., for the liquid vaporizes very slowly unless set on warm radiators or other heated apparatus. The enclosure fumigated must be very tight. This fumigant is purchased as a liquid in tin cans.

#### FUMIGATION WITH MIXTURES OF ETHYLENE OXIDE AND METHYL FORMATE WITH CARBON DIOXIDE

Ethylene oxide-carbon dioxide mixture and the methyl formate-carbon dioxide mixture are two new fumigants that can be used with reasonable safety in fumigating rooms in a house. They produce vapors that are heavier than air and will kill all stages of clothes moths and other household insects. Unfortunately, they are not readily available to the public. They are sold under pressure in steel cylinders, which makes their use difficult except by the professional exterminator. Further information will be furnished by the Department of Agriculture on request.

#### CEDAR-LINED CLOSETS

Closets lined with red cedar wood, as they are ordinarily installed and used in the average American home, are of doubtful value in protecting clothing from moth attack. The experience of house owners and the experimental work of this Department prove that injury from moths often occurs to clothing hung in cedar-lined closets. Usually the closets are not tightly enough constructed to retain the emanations from the cedar wood in sufficient strength to kill the worms. The beautifully contrasting white sapwood is of no value, and the more of it present in the lining the less effective is the closet. As already mentioned (p. 12), the mere odor of cedar does not keep moths from crawling or flying into a closet or from damaging the clothing or furs. Many people enjoy a false sense of security in believing the odor of cedar to be a certain protection.

In justice to red cedar wood it should be stated that closets very tightly constructed with the red heartwood and equipped with doors that close tightly against felt gaskets can be depended upon to protect against moth attack if the doors are clamped shut except when articles are being put in or taken out.

#### CEDAR CHESTS

Well-made chests of red cedar (*Juniperus virginiana*) heartwood, which are in good condition as regards tightness, can be depended on for protection against clothes moths, provided the articles to be placed in them are first thoroughly brushed, combed, or otherwise treated to remove the older clothes-moth larvae. Chests with the sides, ends, and bottoms made of red cedar heartwood at least three-fourths inch thick, and the cover of solid red cedar or of neutral wood lined with red cedar veneer, will kill all the newly hatched or young larvae of clothes moths. On the other hand, in experiments with chests of neutral wood having solid bottoms of  $\frac{3}{4}$ -inch red cedar, but lined on the sides, ends, and tops with  $\frac{1}{20}$ -inch red cedar veneer, clothes moths were able to develop from egg to adult and to cause damage.

The aroma, or the persistent characteristic odor of red cedar, is due to a volatile oil present in the wood to an amount ranging from

1 to 2 percent. The redwood or pure heartwood of red cedar contains from 2 to 4 percent of this oil.<sup>3</sup> As it is the emanation from this volatile oil in the wood that protects clothing, the chests at all times should remain tightly closed except when clothing is being removed or placed in them, and this work should be done as rapidly as possible.

The average trunk in which clothing is stored is not tight enough to keep out moths. Indeed, the unusual tightness of cedar chests, when not cracked or warped, compared with ordinary trunks or other ordinary household receptacles for clothing, gives to such chests a great advantage, aside from their insecticidal value.



FIGURE 19.—Red cedar chests are among the best chests upon the market for the protection of clothing from clothes moths, provided they close tightly and directions for their use given in this bulletin are faithfully followed.

The value of cedar chests (fig. 19) in protecting clothing lies in the fact that they kill the newly hatched or young larvae or worms of clothes moths. Cedar chests cannot be depended on to kill the moths or millers, their eggs, or the worms after they are one-half to full grown or after they are 3 or 4 months old. Neither will the

<sup>3</sup> The principal constituents of the oil are the alcohol cedrol or cedrol camphor, the sesquiterpene alcohol cedrenol, and the sesquiterpene cedrene. The characteristic odor of cedar chests is probably due to the first two compounds mentioned.

chest kill all the pupae or chrysalids. It is because they will not kill these stages of the clothes moth that cedar chests are recommended as protectors of clothing from moth attack only when fabrics put into them are thoroughly cleaned, brushed, or beaten, just before they are placed in the chests. This preliminary treatment before storage, if carefully carried out, with special attention to brushing out pockets, seams, etc., will remove those stages of the moth that the chests will not kill. Remember that if the clothing is not thus brushed, etc., with scrupulous care, some of the older larvae may remain and go into the chests with the clothing and continue to cause damage.

Furs can not always be rid of these older worms by merely brushing or beating. Clothes moth worms often lie hidden next to the skin and are so firmly established by the webbing they spin that they are not dislodged from the fur by brushing or shaking. For this reason furs suspected of being infested should be combed out with a very fine comb, or should be fumigated. If furs are promptly stored as soon as the season for their use is over, however, there is little likelihood that they will contain moth larvae too old to be killed by the chests.

*It is a matter of no practical importance whether cedar chests kill the moths, the eggs, or the pupae, for these stages are not capable of injuring garments. But too much can not be said about the necessity of making sure that clothing going into cedar chests is free from the older larvae or worms; otherwise the owners may sustain losses attributable to carelessness, rather than to the failure of the chest.<sup>4</sup>*

#### CEDAR PRODUCTS

The belief that the mere odor of cedar will keep moths out of trunks, paper bags, boxes, closets, etc., is not based on experience. Red cedar heartwood, as used in good red cedar chests, can be recommended to protect clothing and destroy moths (p. 25). However, the small quantities of cedar oil in certain sprays, in garment bags, storage closets and chests made of corrugated or other cardboards, or in small devices to be hung in closets or rooms, are of no practical value.

Cedar wood oil, cedar leaf oil, or pine oil are often sprayed on paper garment bags, or cardboard storage units in the form of closets or trunks, which are then sold as being "cedarized", as though they were safe storage units because of their odor. As a matter of fact, these storage units are no better from a practical standpoint than if they had not been sprayed with cedar oil or pine oil. The smell of cedar or pine oil does not repel the adult moth or the larva, and weakens and disappears rather soon through evaporation.

Cedar chips and shavings are too apt to be worthless to be recommended.

<sup>4</sup> The foregoing statements regarding the effect of red cedar chests on adults, eggs, and larvae of clothes moths are based on a comprehensive study by the Bureaus of Entomology and Plant Industry of the U. S. Department of Agriculture. The results of this study were reported in Department Bulletin 1051, Red Cedar Chests as Protectors Against Moth Damage, by E. A. Back and Frank Rabak, published in 1922. Copies of this bulletin are no longer available for distribution, but may be consulted in the principal libraries of the United States.

Various plasters and paints, or sheets of cardboard with which to line closets or trunks, have a value too ephemeral to be depended on. Whatever limited value they may possess when honestly made and fresh is soon lost through the evaporation of the cedar oil, and if the surfaces are painted with fresh cedar oil, there is always the danger of the oil oxidizing in the garments and damaging or destroying them.

#### DRY HEAT

Heat is now recognized as an effective agent in killing insects. All fabric pests will be killed in a very short time if rooms can be heated to a temperature of 130° F. and held at that temperature long enough to permit all articles in the room to attain this temperature. Experimental work has demonstrated that even lower temperatures will kill the larvae of clothes moths. All larvae exposed in an incubator to 128°, 120°, and 110° died in 6, 11, and 31 minutes, respectively. At 105°, 20 percent were dead at the end of 31 minutes, while at 110°, 30 percent were dead at the end of 11 minutes. This killing power of heat can be readily utilized by ironing fabrics with very hot irons, or exposing them in superheated rooms before they are put away for the summer.

The rays of the sun in summer can often be used to kill or drive clothes moths from fabrics. Thus clothes-moth eggs on flannel, exposed to the bright sun at 128°, 125°, 120°, and 110° F., were killed at the end of 6, 6, 11, and 31 minutes, respectively. Exposure to the sun at 110° and 105° for 31 and 11 minutes, respectively, did not kill the eggs. Well-grown larvae of clothes moths in garments exposed for several hours to the hottest rays of the sun usually become restless and spin down from the garment. The old-time custom of sunning clothing to kill moths is based on experience.

#### OIL SPRAY

There are on the market a number of sprays, consisting almost entirely of light mineral oils with or without pyrethrum extract in varying quantities, and variously scented. These solutions are used as sprays and to be effective must be brought in actual contact with the insect itself. In other words, they are "contact sprays." Though they are inferior to fumigants in killing out moth infestations, they can be made to kill many moths, including the egg and worm stages, if they are sprayed well into floor cracks, about baseboards, or onto clothing or furniture. They are much more effective when applied with power sprayers. They impart no lasting protection against moths.

#### HOT WATER

Water boils at 212° F. At 115° a person finds water almost too hot to hold the hand in. Fabrics that will not be injured by water can be freed of living clothes-moth eggs and larvae by being dipped for 10 seconds in water heated to 140°. Care, however, must be exercised to have and keep the water at this or a higher temperature. Larvae and eggs in flannel dipped for 10 seconds in water heated only to 122° remained unaffected.

## LAUNDRY SOAP

Clothing washed with a strong solution of neutral laundry soap will be freed from clothes moth larvae and eggs, and if wrapped tightly in paper, as suggested on page 14, as soon as taken from the drying line, will remain free from moth attack.

## GASOLINE

Gasoline sprayed on fabrics kills clothes moths. Dipping clothing in gasoline will kill clothes moths, and articles so dipped and coming direct from the dry-cleaning process may be considered freed, temporarily at least, from moth infestation. Of course, as soon as garments are dried after treatment they are subject to reinfestation. The use of gasoline for the treatment of wearing apparel most subject to moth attack is not advised for the average home because of the dangers and difficulties of application.

## TREATMENT OF CRACKS AND HIDING PLACES WITH GASOLINE, BENZINE, OR KEROSENE

Any effective method of control for clothes moths and other fabric pests must take into consideration the fact that woolen lint, hair, etc., accumulate in floor cracks, behind base boards, and in other such places. In this "dust" carpet beetles and clothes moths can and do subsist and migrate thence to other parts of the house. It is desirable, therefore, to reduce these hiding places as much as possible by the use of crack fillers such as can be purchased at paint shops, and treating them with kerosene, gasoline, or benzene. *Persons are warned against the careless use of these three liquids because of their inflammable nature.* If the premises are fumigated, clothes moths and other fabric pests in floor cracks and similar places are killed, and the more laborious treatment by liquids is not necessary.



# **ORGANIZATION OF THE UNITED STATES DEPARTMENT OF AGRICULTURE WHEN THIS PUBLICATION WAS LAST PRINTED**

---

<i>Secretary of Agriculture</i> .....	HENRY A. WALLACE.
<i>Under Secretary</i> .....	REXFORD G. TUGWELL.
<i>Assistant Secretary</i> .....	M. L. WILSON.
<i>Director of Extension Work</i> .....	C. W. WARBURTON.
<i>Director of Personnel</i> .....	W. W. STOCKBERGER.
<i>Director of Information</i> .....	M. S. EISENHOWER.
<i>Director of Finance</i> .....	W. A. JUMP.
<i>Solicitor</i> .....	SETH THOMAS.
<i>Agricultural Adjustment Administration</i> .....	CHESTER C. DAVIS, <i>Administrator</i> .
<i>Bureau of Agricultural Economics</i> .....	A. G. BLACK, <i>Chief</i> .
<i>Bureau of Agricultural Engineering</i> .....	S. H. MCCRORY, <i>Chief</i> .
<i>Bureau of Animal Industry</i> .....	JOHN R. MOHLER, <i>Chief</i> .
<i>Bureau of Biological Survey</i> .....	J. N. DARLING, <i>Chief</i> .
<i>Bureau of Chemistry and Soils</i> .....	H. G. KNIGHT, <i>Chief</i> .
<i>Office of Cooperative Extension Work</i> .....	C. B. SMITH, <i>Chief</i> .
<i>Bureau of Dairy Industry</i> .....	O. E. REED, <i>Chief</i> .
<i>Bureau of Entomology and Plant Quarantine</i> .....	LEE A. STRONG, <i>Chief</i> .
<i>Office of Experiment Stations</i> .....	JAMES T. JARDINE, <i>Chief</i> .
<i>Food and Drug Administration</i> .....	WALTER G. CAMPBELL, <i>Chief</i> .
<i>Forest Service</i> .....	FERDINAND A. SILCOX, <i>Chief</i> .
<i>Grain Futures Administration</i> .....	J. W. T. DUVEL, <i>Chief</i> .
<i>Bureau of Home Economics</i> .....	LOUISE STANLEY, <i>Chief</i> .
<i>Library</i> .....	CLARIBEL R. BARNETT, <i>Librarian</i> .
<i>Bureau of Plant Industry</i> .....	FREDERICK D. RICHEY, <i>Chief</i> .
<i>Bureau of Public Roads</i> .....	THOMAS H. MACDONALD, <i>Chief</i> .
<i>Weather Bureau</i> .....	WILLIS R. GREGG, <i>Chief</i> .